

Angular momentum of the physical electron

A M Stewart

Department of Theoretical Physics,
Research School of Physical Sciences and Engineering,
The Australian National University,
Canberra, ACT 0200, Australia.

Abstract

The angular momentum of the physical electron, modelled as a Dirac fermion coupled to the electromagnetic field, is found to be $\hbar/2$, the same as that of a bare Dirac fermion and independent of the size of the electric charge.

The coupled Maxwell-Dirac equations

$$(i\gamma^\nu \partial_\nu - \kappa)\psi = \frac{e}{\hbar c} A_\nu \gamma^\nu \psi \quad \text{and} \quad \partial_\sigma F^{\sigma\mu} = 4\pi e \bar{\psi} \gamma^\mu \psi = 4\pi j^\mu / c, \quad (1)$$

where ψ is a field operator and $\kappa = mc/\hbar$, that arise from the Lagrangian density \mathcal{L} of quantum electrodynamics

$$\mathcal{L} / \hbar c = \frac{i}{2} \bar{\psi} \gamma^\sigma \partial_\sigma \psi - \frac{i}{2} (\partial_\sigma \bar{\psi}) \gamma^\sigma \psi - \kappa \bar{\psi} \psi - \frac{e}{\hbar c} A_\sigma \bar{\psi} \gamma^\sigma \psi - \frac{1}{16\pi\hbar c} F^{\alpha\beta} F_{\alpha\beta} \quad (2)$$

imply that the physical electron may be modelled as a composite particle consisting of the bound state of one or more Dirac fermions with bare mass m and charge e and the electromagnetic field [1]. More than one fermion may be necessary to account for the vacuum polarization [2] observed experimentally [3].

Early attempts to calculate physical properties from (1) by perturbation theory led to divergences [4] and more recent attempts to solve (1) as classical equations have met with only limited success [5]. In this note we derive an operator relation for the angular momentum of the physical electron that shows that it has the same spin as a bare Dirac fermion.

The symmetric energy-momentum tensor $T^{\mu\nu}$ that corresponds to the above Lagrangian density is obtained by the usual method [1, 6] to be

$$T^{\mu\nu} = c\hbar \frac{i}{4} (\bar{\psi} \gamma^\mu D^\nu \psi - D^\nu * \bar{\psi} \gamma^\mu \psi + \bar{\psi} \gamma^\nu D^\mu \psi - D^\mu * \bar{\psi} \gamma^\nu \psi) + \frac{1}{4\pi} \left(\frac{\eta^{\mu\nu}}{4} F_{\alpha\beta} F^{\alpha\beta} + F^{\mu\sigma} F_\sigma^\nu \right) \quad (3)$$

where $D^\mu = \partial^\mu + \frac{ie}{\hbar c} A^\mu$ is the gauge covariant derivative. The electromagnetic part of the tensor has been made symmetric and gauge invariant by the method of Belinfante [1, 7].

From (3) the angular momentum operator \mathbf{J} of the coupled system is found to be

$$\mathbf{J} = -i\hbar \int d^3x \mathbf{x} \times \psi^\dagger (\nabla - \frac{ie}{\hbar c} \mathbf{A}) \psi + \frac{\hbar}{2} \int d^3x \psi^\dagger \boldsymbol{\Sigma} \psi + \frac{1}{4\pi c} \int d^3x \mathbf{x} \times (\mathbf{E} \times \mathbf{B}) \quad (4)$$

where the four-component spin operator $\boldsymbol{\Sigma}$ is

$$\boldsymbol{\Sigma} = \begin{bmatrix} \boldsymbol{\sigma} & 0 \\ 0 & \boldsymbol{\sigma} \end{bmatrix}, \quad (5)$$

$\boldsymbol{\sigma}$ being the three Pauli matrices [1]. The calculation of the Dirac parts of (4) from (3) is given by Ohanian [8]. The last term is the angular momentum of the electromagnetic field. The expression for \mathbf{J} is gauge invariant.

The angular momentum of the electromagnetic field has been examined recently [9, 10, 11, 12, 13], see also [14]. It can be separated into two parts, a radiative part and a bound part. The radiative part may in turn be decomposed into three terms, a spin term, an orbital term and a boundary term. The spin term and orbital term have been shown to give the standard expressions for paraxial waves [13]. The spin term and the boundary term have been shown to resolve the paradox involving the angular momentum of plane waves of electromagnetic radiation [11]. However, because an electron in uniform motion does not radiate, only the bound part needs to be considered in the present context. The bound part is [12]

$$\mathbf{J}_b = \frac{1}{c} \int d^3x \rho(\mathbf{x}) \mathbf{x} \times \mathbf{A}_t(\mathbf{x}) \quad (6)$$

where \mathbf{A}_t is the vector potential in the transverse or Coulomb gauge ($\text{div} \mathbf{A} = 0$) and ρ is the charge density, in the present case $e\psi^\dagger(\mathbf{x})\psi(\mathbf{x})$. The transverse vector potential can be expressed explicitly either in terms of the instantaneous magnetic field [15, 16] or in terms of the retarded current [17]. The exact forms need not concern us.

If we choose to work in the Coulomb gauge with $\mathbf{A} = \mathbf{A}_t$ we find that the two terms that contain the vector potential cancel and (4) simplifies to give

$$\mathbf{J} = -i\hbar \int d^3x \mathbf{x} \times \psi^\dagger \nabla \psi + \frac{\hbar}{2} \int d^3x \psi^\dagger \boldsymbol{\Sigma} \psi \quad (7)$$

Although (7) gives the angular momentum operator of the full interacting system it has the same form as for a non-interacting Dirac fermion. Accordingly, the spin of the physical electron, like that of the Dirac fermion, has a magnitude of $\hbar/2$, independent of the size of the electronic charge. Although the electromagnetic field has an angular momentum given by the last term of

(4) [18], it is exactly cancelled by the term in (4) that exhibits the vector potential explicitly. Since (4) is gauge invariant the argument holds in any gauge.

A similar result holds for the linear momentum. The result for angular momentum obtained here would appear to demonstrate the futility of any attempt to relate the quantization of angular momentum to the quantization of electric charge.

References

- [1] Greiner W and Reinhardt J 1996 *Field Quantization* (Berlin: Springer)
- [2] Schwinger J 1951 Gauge Invariance and Vacuum Polarisation *Phys. Rev.* **82** 664-79
- [3] Levine I, Kohtick D, Howell B, *et al.* 1997 Measurement of the Electromagnetic Coupling at Large Momentum Transfers *Phys. Rev. Lett.* **78** 424-7
- [4] Weisskopf V F 1939 On the self-energy of the electromagnetic field of the electron *Phys. Rev.* **56** 72-85
- [5] Radford C and Booth H 1999 Magnetic monopoles, electric neutrality and the static Maxwell-Dirac equations *J. Phys. A: Math. Gen.* **32** 5807-22
- [6] Gupta S N 1977 *Quantum Electrodynamics* (New York: Gordon and Breach)
- [7] Belinfante F J 1939 On the Spin Angular Momentum of Mesons *Physica* **6** 887-98
- [8] Ohanian H C 1986 What is spin? *Am. J. Phys.* **54** 500-5
- [9] Stewart A M 2005 Angular momentum of light *J. Mod. Opt.* **52** 1145-54
- [10] Stewart A M 2005 *General expression for the orbital and spin components of the angular momentum of light*. Proceedings of the 16th National Biennial Congress of the Australian Institute of Physics, Canberra, (Canberra: Australian Institute of Physics)
- [11] Stewart A M 2005 Angular momentum of the electromagnetic field: the plane wave paradox resolved *Eur. J. Phys.* **26** 635-41
- [12] Stewart A M 2005 Equivalence of two mathematical forms for the bound angular momentum of the electromagnetic field *J. Mod. Opt.* **52** 2695-8
- [13] Stewart A M 2006 Derivation of the paraxial form of the angular momentum of the electromagnetic field from the general form *J. Mod. Opt.* **53** 1947-52
- [14] Cohen-Tannoudji C, Dupont-Roc J and Gilbert G 1989 *Photons and Atoms* (New York: Wiley)
- [15] Stewart A M 2003 Vector potential of the Coulomb gauge *Eur. J. Phys.* **24** 519-24
- [16] Stewart A M 2004 Reply to Comments on 'Vector potential of the Coulomb gauge' *Eur. J. Phys.* **25** L29-L30
- [17] Jackson J D 2002 From Lorenz to Coulomb and other explicit gauge transformations *Am. J. Phys.* **70** 917-28
- [18] Highbe J 1988 Angular momentum of the field of an electron *Am. J. Phys.* **56** 378-9